

# QUERTANIN®

Stave wood quality ellagic tannins, extracted from oak heartwood, using LAFFORT®'s Instant Dissolving Process (IDP).  
For post-vinification use in red, rosé and white wine.

*Qualified for the elaboration of products for direct human consumption in the field of the regulated use in oenology.  
In accordance with the current EU regulation n° 2019/934.*

## SPECIFICATIONS AND OENOLOGICAL APPLICATIONS


QUERTANIN® is a preparation of ellagitannins with a strong capacity for complex formation, used in vinification or during ageing, with antioxidant and antioxidase properties (inhibition of laccase) and high reactivity with proteins (facilitates clarification).

- Regulates oxidation-reduction phenomena during maturation in barrels or during micro-oxygenation.
- Treats wines showing reductive characteristics.
- Creates an environment rich in ellagic tannins in used barrels, similar to that of new barrels.


## EXPERIMENTAL RESULT

- Tannin content extracted from the oak is lower in used barrels. The ellagitannin protective effect is decreased and the wine becomes subject to premature oxidation. Adding QUERTANIN® permits the re-creation of the buffer qualities provided by tannins extracted from new barrels, and protects the wine from oxidation phenomena.
- QUERTANIN® permits a significant decrease in reductive characters, such as light mercaptans, guaranteeing improved preservation of aromatic freshness.

ELLAGITANNIN CONTENT IN BARRELS



ETHANETHIOL CONTENT (%)



**LAFFORT**  
*l'œnologie par nature*

## PHYSICAL CHARACTERISTICS

---

Aspect .....	granulated	Colour .....	dark brown
Solubility .....	complete		

## CHEMICAL ANALYSIS

---

Tannins (%) .....	> 65	Arsenic (ppm) .....	< 3
Humidity (%) .....	< 10	Iron (ppm) .....	< 50
Ash (%) .....	< 5	Lead (ppm) .....	< 2
Insoluble substances (%) .....	< 5	Mercury (ppm) .....	< 0.5
		Cadmium (ppm) .....	< 0.5

## PROTOCOL FOR USE

### DOSAGE

*It is specified in the Oenological Codex that tannins "must not alter the recognised organoleptic characteristics of the wines". The dosage rates will therefore vary in function of the wine matrix, and shall be determined after trials.*

- Clarification additive: 10 - 30 g/hL (100 - 300 ppm).
- Inhibition of laccase activity: 30 – 80 g/hL (300 - 800 ppm).
- Prevention of oxidation in white and rosé wines: 2 - 5 g/hL (20 - 50 ppm).
- Red wine ageing: 5 - 20 g/hL (50 - 200 ppm).

### IMPLEMENTATION

Thanks to the IDP process, QUERTANIN® can be sprinkled directly onto the wine during homogenisation or a pump-over (1 - 10%).

After QUERTANIN® has been added, it is recommended to carry out normal rackings until bottling preparation.

It is recommended to add QUERTANIN® at least two weeks before bottling.

*For white wines stabilised with CMC (tartaric stabilisation), the addition of tannins towards the end of ageing may affect the colloidal stability; it is recommended to perform a protein stability test again systematically after addition.*

## STORAGE RECOMMENDATION

- 
- Store above ground level in a dry area not liable to impart odours. Ensuring stock is kept at a moderate temperature, in its original, unopened packaging.
  - Optimal date of use: 5 years.

## PACKAGING

---

500 g bag - 10 kg box.

